



Unexpected interactions between gold and N-morpholino-sulfonates

Nielsen, Frederick Stappen; Engelbrekt, Christian; Elliot, Samuel Gilbert; Junor, Glen; Sørensen, Kasper; Jensen, Knud J.; Clausen, Mads Hartvig; Zhang, Jingdong

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Nielsen, F. S., Engelbrekt, C., Elliot, S. G., Junor, G., Sørensen, K., Jensen, K. J., Clausen, M. H., & Zhang, J. (2017). *Unexpected interactions between gold and N-morpholino-sulfonates*. Abstract from Forth EuCheMS Inorganic Chemistry Conference (EICC-4), Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Unexpected interactions between gold and N-morpholino-sulfonates

Frederick Stappen¹, Christian Engelbrekt¹, Samuel Eliot¹, Glen Junor², Kasper Sørensen³, Knud J. Jensen³, Mads H. Clausen¹ and Jingdong Zhang¹.

¹Department of Chemistry, Technical University of Denmark, ²Department of Chemistry, University of California San Diego, ³Department of Chemistry, University of Copenhagen. fcksn@kemi.dtu.dk

Nanoporous gold (NPG) has a high surface area and excellent conductivity. It is an ideal supporting material for the electrocatalysis, e.g. in fuel cell applications. NPG is traditionally produced by etching a gold/silver alloy. This method has significant drawbacks, such as the introduction of silver into your NPG, and its multi-step fabrication. A method has been discovered for producing NPG as a thin film chemically. This bottom-up approach entails reduction of Au³⁺ precursor using morpholinoethanesulfonic acid (MES). This produces a thin and highly porous gold film at the air-water interface¹ (for details, see poster by Mikkel Christiansen).

This chemical reaction is far more complex than first expected and bi-products, intermediates and reaction mechanisms are the focus of the present work.

The chemical reaction and its products have been examined using state-of-the-art nuclear magnetic resonance (NMR), ultraviolet-visible spectroscopy (UV-vis), fluorescence spectroscopy, density functional theory (DFT), mass spectrometry (MS) and Raman spectroscopy. The results illustrate a complex chain of reactions resulting in gold nanoparticles, NPG, and several previously unidentified Au-complexes. N-NMR presented three different environments for the N-atom in MES, while the UV-vis results points towards some interesting gold complexes. MS identifies several distinct molecular entities demonstrating the reactivity of MES and Au, and fluorescence spectroscopy suggests the formation of polynuclear Au complexes as previously reported where Au nuclei are bridged by C=N functionalities in small organic molecules². This adds up to a complicated reaction mechanism involving some interesting Au^{3+/1+} complexes, deprotonation and oxidation of MES and the formation of molecules that show UV-vis absorbance and fluorescence.




Figure 1 – (left), UV-vis spectra of AuCl_x(H₂O)_y complexes corresponding to (right) those found after the reaction between gold and MES.

- (1) Engelbrekt, C.; Wagner, M.; Christiansen, M. U.-B.; Christensen, H. E. M.; Qian, X.; Ulstrup, J.; Zhao, C.; Zhang, J. Side Effect of Good's Buffers on Optical Properties of Gold Nanoparticle Solutions. *ChemElectroChem* **2016**, 2015–2016.
- (2) Fung, E. Y.; Olmstead, M. M.; Vickery, J. C.; Balch, A. L. Glowing Gold Rings: Solvoluminescence from Planar trigold(I) Complexes. *Coord. Chem. Rev.* **1998**, 171, 151–159.